

R. Van Malderen (1,5), E. Pottiaux (2,5), H. Brenot (3,5), S. Beirle (4), T. Wagner (4), C. Hermans (3), M. De Mazière (3), H. De Backer (1) and C. Bruyninx (2)

(1) Royal Meteorological Institute of Belgium, (2) Royal Observatory of Belgium, (3) Belgian Institute for Space Aeronomy, (4) Max Planck Institute for Chemistry (MPI-C), Mainz, Germany, (5) Solar-Terrestrial Centre of Excellence

Mail to: Roeland.VanMalderen@meteo.be or Eric.Pottiaux@observatoire.be

GOAL


Water vapour plays a dominant role in the climate change debate. However, observing water vapour for a climatological time period in a consistent and homogeneous manner is a challenging task. To this end, water vapour estimations derived from ground-based observations of **Global Navigation Satellite System (GNSS)** receiver networks such as the International GNSS Service (IGS) network are very promising, with continuous observations spanning over the last 15+ years. In addition, the **AEROSOL ROBOTIC NETWORK (AERONET)** also provides long-term and continuous ground-based observations of the total water vapour content performed with standardized and well-calibrated sun photometers.

The present study aims to assess the applicability of either datasets for **water vapour time series analysis**. Therefore, we compare the Integrated Water Vapour (IWV) measurements retrieved from these two techniques, focusing on a selection of almost 30 sites worldwide. Furthermore, we compare the GPS and sun photometer IWV values with simultaneous and co-located radiosonde and satellite-based IWV measurements (**GOME(2)** and **SCIAMACHY**) and we investigate the geographical dependency of the properties of the IWV scatter plots between all these different instruments.

1. INSTRUMENTS AND DATASETS

INSTRUMENTS:

- 2 Ground-Based Instruments
- 1 In-Situ Instrument
- 3 Satellite-Based Instruments



Technique	Spatial Coverage	Temporal Resolution	Time Span	Tech. Costs	All Weather / All Directions	By Product of An Analysis
GPS	± 350 IGS sites	every 5 minutes	1995-now	low	Yes / Yes	Yes
Radiosonde	± 1500 sites	on average twice/day	1950s-now	low to moderate	Yes / Vertical Profile	No
CIMEL Sun Photometers	± 300 sites	± 15 min, depending on weather conditions	1993-now	moderate	clear sky only / solar direction needed	No, but focus on aerosol properties retrieval
GOME(2)/SCIAMACHY	Global	maximum twice/day	1996-now	very high	only if (almost) cloud free/nadir	No

Table 1 : Pros & cons per technique.




Fig. 1: Map of the selected sites that host at least 2 of the considered instruments.

Within a maximum separating distance of 30 km, 28 co-locations are found worldwide between IGS GPS sites and AERONET CIMEL sun photometer locations (see Fig. 1). Additionally, we looked for radiosonde launches and GOME(2) & SCIAMACHY crossings at those selected sites. The IWV data sets from the different instruments are retrieved as follows:

- **GPS:** GPS-based Zenith Total Delay (ZTD) from the IGS Final (re)processing (Byun and Bar-Sever [2009,2010]) is converted into IWV by using surface measurements of temperature and pressure, gathered at synoptic stations at a horizontal distance of maximum 50 km from the GPS station (more details in e.g. Bevis et al. [1992], Saastamoinen [1972], Askne and Nordius [1987] and Davis et al. [1985]).
- **CIMEL:** IWV is obtained by measuring the (direct) sun radiance at a 940nm channel (centred on the 946nm water vapour absorption line).
- **Radiosondes:** IWV is calculated through integration of the vertical profiles of temperature and relative humidity.
- **GOME/SCIAMACHY/GOME-2 (=GOMESCIA):** IWV is retrieved by applying the so-called Air Mass Corrected Differential Optical Absorption Spectroscopy method to nadir measurements around 700nm.

The advantages and disadvantages of each technique are summarized in Table 1.

2. TECHNIQUES INTERCOMPARISON

INSTRUMENT COLOCATION: FOCUS ON BRUSSELS

As a first step, this study focused on Uccle, Brussels, Belgium (50°48'N, 4°21'E, 100m asl) presenting the following advantages:

- The different ground-based and in-situ instruments and the automatic weather station (time resolution: 10min) are really located at the same site, so that the horizontal and vertical separation of the different devices is not an issue.
- All techniques are available for this site.
- We dispose of the metadata of the different instruments, so that we are aware of any instrumental change that might give rise to an inhomogeneity of the instrument's data series.
- The availability of auxiliary weather data is a major advantage.

From Fig. 2, we note:

- The different instruments have different observation periods.
- We have 2 radiosonde types: Vaisala's RS80 and RS90/RS92 (=RS9x).
- The GPS IGS IWV is candidate for reference device because of data every 10min (since 1999*), only minor data gaps, homogeneous data (re)-processing by IGS.

* We dispose of weather data with 10 minutes of time resolution only since 1999.




Fig. 2: Overview of all IWV data available at Uccle, Brussels.

EXPLOITATION OF THE IWV DATASETS @ BRUSSELS

We constructed scatter plots of simultaneous ($\Delta t = 10$ min for CIMEL, $\Delta t = 30$ min for RS and GOMESCIA) IWV measurements between the different devices (using the GPS as reference, see Fig. 3). These plots show that:

- The mean bias between the different techniques varies between **-0.6 mm (GOMESCIA) to 0.6 mm (RS9x)**.
- The best correlation and lowest dispersion of the data points are reached for the **CIMEL vs. GPS** comparison.
- Vaisala's state-of-the-art radiosonde type (RS9x) compares better w.r.t. GPS data than the preceding RS80 type.
- The slopes of regression lines w.r.t. GPS are closer to 1 for other all-weather devices (RS) than for instruments demanding a partly clear sky (CIMEL, GOMESCIA). A small study incorporating the available cloud cover data demonstrated that the presence of clouds leads to higher IWV values, especially for GPS observations, compared to the simultaneous CIMEL measurements.




Fig. 3: Scatter plots of simultaneous IWV measurements of the different instruments with the GPS device.

WORLD-WIDE EXPLOITATION OF IWV DATASETS

In a second step, we extended our study worldwide. We created scatter plots similar to Fig. 3 for the selected 28 sites for which we found instrumental co-location. Results are summarised in Figs. 4 and 5 and show that:

- The **CIMEL instrument compares best with the GPS** technique for the IWV measurements (best correlation, lowest scatter).
- The **regression slopes** are for almost all instrument comparisons at all stations **smaller than 1**.
- At sites where different CIMELs can be compared with one IGS GPS station (e.g. BRMU, NISU, TLSE, BUCU, VENE, OBE2, OPMT), significant differences exist between the regression slopes of the respective scatter plots → geographical dependency or remaining CIMEL calibration issues?
- There is **neither latitudinal nor longitudinal dependency** of the scatter plots properties.




Fig. 4: Column bar plots of scatter plot properties (count N, bias, R^2 and regression slope) of the different instruments versus GPS for the selected sites worldwide. Sites are ordered with increasing latitude. The error bars represent the RMS (bias) and the standard deviation (regression slope).




Fig. 5: Column bar plots of scatter plot properties (count N, bias, R^2 and regression slope) of the different instruments versus GPS averaged over all stations included in the inter-technique comparison. Error bars: see Fig. 4.

3. CONCLUSIONS AND PERSPECTIVES

• **CIMEL sun photometers and GPS** are very valuable ground-based techniques and therefore good candidates to evaluate/calibrate the long-term IWV datasets provided by satellite devices: they correlate very well (their mean R^2 is 0.984) and **typically agree at the level of $0.3 \text{ mm} \pm 1.5 \text{ mm}$ of IWV**.

• **Influence of clouds:** for large IWV values, the GPS instrument measures higher amounts of IWV than the CIMEL does. This can at least partly be explained by the observation bias of the CIMEL instrument: it requires a clear sky in the direction of the sun. But the larger the IWV values, the higher the probability to have clouds, which contribute directly to the GPS observations, but not to the CIMEL IWV observations.

• Both ground-based techniques are very **promising to build up long time series for climate applications**, as long as the data homogeneity can be guaranteed. For the CIMEL photometers belonging to the AERONET, a regular calibration of the instrument is required. IGS GPS data were (re)processed homogeneously from 1994 on to mid-April 2011.

• As a first example, we evaluated the **GOMESCIA** satellite IWV dataset. The GOMESCIA IWV measurements are susceptible to a similar observation bias as the CIMEL (almost cloud free skies are needed), which is also reflected in the low mean value of the regression line slope. For these satellite data, the **largest (but apparently random) geographical variability** of the IWV measurements relative to the co-located GPS observations is obtained.

• At this point of our research, there is **no clear geographical pattern** (e.g. related to the climate type) in the inter-technique comparisons at the selected sites worldwide.

• This research has now been **extended to other (e.g. IR) satellite devices** measuring IWV (AIRS) and is submitted to ACP.

REFERENCES AND ACKNOWLEDGEMENTS

This research has been carried out in the framework of the Solar-Terrestrial Centre of Excellence (STCE). We are grateful to all colleagues and data providers below:



REFERENCES:

- Askne, J. and Nordius, H. (1987). Estimation of tropospheric delay for microwaves from surface weather data. *Radio Science*, 22:379-386
- Bevis, M., S. Businger, T. A. Herring, C. Rocken, R. A. Anthes, and R. Ware (1992). GPS meteorology: Remote sensing of atmospheric water vapor using the Global Positioning System. *J. Geophys. Res.*, 97(D14), 15787-15801.
- Byun, S. H., and Bar-Sever, Y. E. (2009). A new type of troposphere zenith path delay product of the international GNSS service. *Journal of Geodesy*, 83(3-4), pp.1-7.
- Byun, S. H., and Bar-Sever, Y. E. (2010). The Re-Analysis of the IGS Tropospheric Product. Available at: http://acc.igs.org/trop/trop-repro_IGSW10.pdf.
- Davis, J. L., Herring, T. A., Shapiro, I. I., Rogers, A. E. E., and Elgered, G. (1985). Geodesy by radio interferometry: effects of atmospheric modelling errors on estimates of baseline length. *Radio Science*, 20(6):1593-1607.
- Saastamoinen, J. (1972). Atmospheric correction for troposphere and stratosphere in radio ranging of satellites. *The Use of Artificial Satellites for Geodesy*, Third International Symposium on The Use of Artificial Satellites for Geodesy

